CARBOPLATIN INJECTION

Rx only

Carboplatin injection should be administered under the supervision of a qualified physicial experienced in the use of cancer chemotherapeutic agents. Appropriate management of therapy and complications is possible only when adequate treatment facilities are readily

Bone marrow suppression is dose related and may be severe, resulting in infection and/ or bleeding. Anemia may be cumulative and may require transfusion support. Vomiting is another frequent drug-related side effect.

Anaphylactic-like reactions to carboplatin have been reported and may occur within minutes of carboplatin injection administration. Epinephrine, corticosteroids, and antihistamines have been employed to alleviate symptoms.

Carboplatin injection is supplied as a sterile, pyrogen-free solution available in 10 mg per mL $\,$ multiple-dose vials containing 50 mg, 150 mg and 450 mg of carboplatin for administration by intravenous infusion. Each mL contains: carboplatin 10 mg and water for injection to volume. Carboplatin is a platinum coordination compound. The chemical name for carboplatin is Platinum, diammine[1,1-cyclobutanedicarboxylato(2-)-O,O]-,(SP-4-2), and carboplatin has

Carboplatin, USP is a white to off-white solid. It is soluble in water, very slightly soluble in acetone and in alcohol. It is soluble in dimethylacetamide and dimethylformamide, and the pH of a 1% solution is 5 to 7.

C₆H₁₂N₂O₄Pt M.W. 371.25

CLINICAL PHARMACOLOGY

DNA-protein cross-links. This effect is apparently cell-cycle nonspecific. The aquation of carboplatin, which is thought to produce the active species, occurs at a slower rate than in the case of cisplatin. Despite this difference, it appears that both carboplatin and cisplatin induce equal numbers of drug-DNA cross-links, causing equivalent lesions and biological effects. The differences in potencies for carboplatin and cisplatin appear to be directly related to the difference in aquation rates.

In patients with creatinine clearances of about 60 mL/min or greater, plasma levels of intact carboplatin decay in a biphasic manner after a 30-minute intravenous infusion of 300 mg/m² to 500 mg/m² of carboplatin. The initial plasma half-life (alpha) was found to be 1.1 to 2 hours (n = 6), and the postdistribution plasma half-life (beta) was found to be 2.6 to 5.9 hours (n = 6). The total body clearance, apparent volume of distribution and mean residence time for carboplatin are 4.4 L/hour, 16 L and 3.5 hours, respectively. The C $_{max}$ values and areas under the plasma concentration versus time curves from 0 to infinity (AUC inf) increase linearly with dose, although the increase was slightly more than dose proportional. Carboplatin, therefore, $% \left(1\right) =\left(1\right) \left(1\right) \left$ exhibits linear pharmacokinetics over the dosing range studied (300 mg/m² to 500 mg/m²).

Carboplatin is not bound to plasma proteins. No significant quantities of protein-free, ultrafilterable platinum-containing species other than carboplatin are present in plasma. However, platinum from carboplatin becomes irreversibly bound to plasma proteins and is slowly eliminated with a minimum half-life of 5 days.

The major route of elimination of carboplatin is renal excretion. Patients with creatinine clearances of approximately 60 mL/min or greater excrete 65% of the dose in the urine within 12 hours and 71% of the dose within 24 hours. All of the platinum in the 24-hour urine is present as carboplatin. Only 3 to 5% of the administered platinum is excreted in the urine between 24 and 96 hours. There are insufficient data to determine whether biliary excretion

In patients with creatinine clearances below 60 mL/min, the total body and renal clearances of carboplatin decrease as the creatinine clearance decreases. Carboplatin dosages should therefore be reduced in these patients (see **DOSAGE AND ADMINISTRATION**).

The primary determinant of carboplatin injection clearance is glomerular filtration rate (GFR) and this parameter of renal function is often decreased in elderly patients. Dosing nulas incorporating estimates of GFR (see DOSAGE AND ADMINISTRATION) to provide predictable carboplatin injection plasma AUCs should be used in elderly patients to minimize

CLINICAL STUDIES

Use with Cyclophosphamide for Initial Treatment of Ovarian Cancer

In two prospectively randomized, controlled studies conducted by the National Cancer Institute of Canada, Clinical Trials Group (NCIC) and the Southwest Oncology Group (SWOG), 789 chemotherapy naive patients with advanced ovarian cancer were treated with carboplatin or cisplatin, both in combination with cyclophosphamide every 28 days for 6 courses before surgical reevaluation. The following results were obtained from both studies:

(Carboplatin-Cisplatin)

oniparative Enicacy			
Overview of Pivotal Trials			
	NCIC	SWOG	
Number of patients randomized	447	342	
Median age (years)	60	62	
Dose of cisplatin	75 mg/m ²	100 mg/m ²	
Dose of carboplatin	300 mg/m ²	300 mg/m ²	
Dose of cyclophosphamide	600 mg/m ²	600 mg/m ²	
Residual tumor < 2 cm (number of patients)	39% (174/447)	14% (49/342)	

Clinical Response in Measurable Disease Patients		
	NCIC	SWOG
Carboplatin (number of patients)	60% (48/80)	58% (48/83)
Cisplatin (number of patients)	58% (49/85)	43% (33/76)
95% CI of difference	(-13.9%, 18.6%)	(-2.3%, 31.1%)

Pathologic Complete Response

	NCIC	SWOG
Carboplatin (number of patients)	11% (24/224)	10% (17/171)
Cisplatin (number of patients)	15% (33/223)	10% (17/171)
95% CI of difference	(-10.7%, 2.5%)	(-6.9%, 6.9%)
(Carboplatin–Cisplatin)		

* 114 Carboplatin and 109 Cisplatin patients did not undergo second look surgery in NCIC study. 90 Carboplatin and 106 Cisplatin patients did not undergo second look surgery in SWOG study.

Progression-Free Survival (PFS)

	NCIC	SWOG
Median		
Carboplatin	59 weeks	49 weeks
Cisplatin	61 weeks	47 weeks
2 year PFS*		
Carboplatin	31%	21%
Cisplatin	31%	21%
95% CI of difference	(-9.3, 8.7)	(-9, 9.4)
(Carboplatin–Cisplatin)		
3 year PFS*		
Carboplatin	19%	8%
Cisplatin	23%	14%
95% CI of difference	(-11.5, 4.5)	(-14.1, 0.3)
(Carboplatin–Cisplatin)		
Hazard Ratio [†]	1.1	1.02
95% CI	(0.89, 1.35)	(0.81, 1.29)
(Carboplatin–Cisplatin)		
* Kaplan-Meier Estimates		

Survival		
	NCIC	SWOG
Median		
Carboplatin	110 weeks	86 weeks
Cisplatin	99 weeks	79 weeks
2 year Survival*		
Carboplatin	51.9%	40.2%
Cisplatin	48.4%	39%
95% CI of difference	(-6.2, 13.2)	(-9.8, 12.2)
Carboplatin–Cisplatin)		
3 year Survival*		
Carboplatin	34.6%	18.3%
Cisplatin	33.1%	24.9%
95% CI of difference	(-7.7, 10.7)	(-15.9, 2.7)
Carboplatin–Cisplatin)		
lazard Ratio [†]	0.98	1.01
95% CI	(0.78, 1.23)	(0.78, 1.3)
Carboplatin–Cisplatin)		
Mojor Entimatos		

Analysis adjusted for factors found to be of prognostic significance were consistent with unadjusted

The pattern of toxicity exerted by the carboplatin-containing regimen was significantly different from that of the cisplatin-containing combinations. Differences between the two studies may be explained by different cisplatin dosages and by different supportive care.

The carboplatin-containing regimen induced significantly more thrombocytopenia and, in one study, significantly more leukopenia and more need for transfusional support. The cisplating containing regimen produced significantly more anemia in one study. However, no significant differences occurred in incidences of infections and hemorrhagic episodes.

Non-hematologic toxicities (emesis, neurotoxicity, ototoxicity, renal toxicity, hypomagnesemia and alopecia) were significantly more frequent in the cisplatin-containing arms.

ADVERSE EXPERIENCES IN PATIENTS WITH OVARIAN CANCER NCIC STUDY

		Carboplatin	Cisplatin	
		Arm	Arm	
		Percent*	Percent*	P-Values [†]
one Marrow				
Thrombocytopenia	< 100,000/mm ³	70	29	< 0.001
	< 50,000/mm ³	41	6	< 0.001
Neutropenia	< 2000 cells/mm ³	97	96	ns
	< 1000 cells/mm ³	81	79	ns
_eukopenia	< 4000 cells/mm ³	98	97	ns
	< 2000 cells/mm ³	68	52	0.001
Anemia	< 11 g/dL	91	91	ns
	< 8 g/dL	18	12	ns
nfections		14	12	ns
Bleeding		10	4	ns
ransfusions -		42	31	0.018
astrointestinal				
Nausea and vomiting	ng	93	98	0.01
/omiting		84	97	< 0.001
Other GI side effect	s	50	62	0.013
eurologic				
Peripheral neuropa	thies	16	42	< 0.001
Ototoxicity		13	33	< 0.001
Other sensory side	effects	6	10	ns
Central neurotoxicit	y	28	40	0.009
enal	•			
Serum creatinine el	evations	5	13	0.006
Blood urea elevatio	ns	17	31	< 0.001

ADVERSE EXPERIENCES IN PATIENTS WITH OVARIAN CANCER NCIC STUDY (cont'd)

	Arm	Arm	
	Percent*	Percent*	P-Values [†]
Hepatic			
Bilirubin elevations	5	3	ns
SGOT elevations	17	13	ns
Alkaline phosphatase elevations	-	-	-
Electrolytes loss			
Sodium	10	20	0.005
Potassium	16	22	ns
Calcium	16	19	ns
Magnesium	63	88	< 0.001
Other side effects			
Pain	36	37	ns
Asthenia	40	33	ns
Cardiovascular	15	19	ns
Respiratory	8	9	ns
Allergic	12	9	ns
Genitourinary	10	10	ns
Alopecia [‡]	50	62	0.017
Mucocitic	10	٥	ne

/alues are in percent of evaluable patients

ADVERSE EXPERIENCES IN PATIENTS WITH OVARIAN CANCER SWOG STUDY

		Carbopiatiii	Cispiatifi	
		Arm	Arm	
		Percent*	Percent*	P-Values [†]
Bone Marrow				
Thrombocytopenia	< 100,000/mm ³	59	35	< 0.001
	< 50,000/mm ³	22	11	0.006
Neutropenia	< 2,000 cells/mm ³	95	97	ns
	< 1,000 cells/mm ³	84	78	ns
Leukopenia	<4,000cells/mm ³	97	97	ns
	<2,000cells/mm ³	76	67	ns
Anemia	< 11 g/dL	88	87	ns
	< 8 g/dL	8	24	< 0.001
Infections		18	21	ns
Bleeding		6	4	ns
Transfusions		25	33	ns
Gastrointestinal				
Nausea and vomiting		94	96	ns
Vomiting		82	91	0.007
Other GI side effects		40	48	ns
Neurologic				
Peripheral neuropathies		13	28	0.001
Ototoxicity		12	30	< 0.001
Other sensory side	;	4	6	ns
effects				
Central neurotoxicity		23	29	ns
Renal				
Serum creatinine	9	7	38	< 0.001
elevations				
Blood urea elevations		-	-	-
Hepatic				
Bilirubin elevations		5	3	ns
SGOT elevations		23	16	ns
Alkaline phospha	tase	29	20	ns
elevations				
Electrolytes loss				

ADVERSE EXPERIENCES IN PATIENTS WITH OVARIAN CANCER SWOG STUDY (cont'd)

	Carboplatin	Cisplatin	
	Arm	Arm	
	Percent*	Percent*	P-Values [†]
Potassium	-	-	-
Calcium	-	-	-
Magnesium	58	77	< 0.001
Other side effects			
Pain	54	52	ns
Asthenia	43	46	ns
Cardiovascular	23	30	ns
Respiratory	12	11	ns
Allergic	10	11	ns
Genitourinary	11	13	ns
Alopecia [‡]	43	57	0.009
Mucositis	6	11	ns

ns = not significant, p > 0.05

* May have been affected by cyclophosphamide dosage delivered

Use as a Single Agent for Secondary Treatment of Advanced Ovarian Cancer

In two prospective, randomized controlled studies in patients with advanced ovarian cancer previously treated with chemotherapy, carboplatin achieved 6 clinical complete responses in 47 patients. The duration of these responses ranged from 45 to 71+ weeks.

INDICATIONS

Initial Treatment of Advanced Ovarian Carcinoma

Carboplatin injection is indicated for the initial treatment of advanced ovarian carcinoma in established combination with other approved chemotherapeutic agents. One established combination regimen consists of carboplatin and cyclophosphamide. Two randomized controlled studies conducted by the NCIC and SWOG with carboplatin versus cisplatin, both in combination with cyclophosphamide, have demonstrated equivalent overall survival between the two groups (see CLINICAL STUDIES).

There is limited statistical power to demonstrate equivalence in overall pathologic complete response rates and long-term survival (≥ 3 years) because of the small number of patients with these outcomes: the small number of patients with residual tumor < 2 cm after initial surgery also limits the statistical power to demonstrate equivalence in this subgroup.

Secondary Treatment of Advanced Ovarian Carcinoma

Carboplatin injection is indicated for the palliative treatment of patients with ovarian carcinoma recurrent after prior chemotherapy, including patients who have been previously treated with cisplatin.

Within the group of patients previously treated with cisplatin, those who have developed progressive disease while receiving cisplatin therapy may have a decreased response rate.

CONTRAINDICATIONS

Carboplatin injection is contraindicated in patients with a history of severe allergic reactions to cisplatin or other platinum-containing compounds.

Carboplatin injection should not be employed in patients with severe bone marrow depression

WARNINGS

Bone marrow suppression (leukopenia, neutropenia, and thrombocytopenia) is dosedependent and is also the dose-limiting toxicity. Peripheral blood counts should be frequently monitored during carboplatin injection treatment and, when appropriate, until recovery is achieved. Median nadir occurs at day 21 in patients receiving single agent carboplatin. In general, single intermittent courses of carboplatin should not be repeated until leukocyte, neutrophil, and platelet counts have recovered.

Since anemia is cumulative, transfusions may be needed during treatment with carboplatin, particularly in patients receiving prolonged therapy.

Bone marrow suppression is increased in patients who have received prior therapy, especially regimens including cisplatin. Marrow suppression is also increased in patients with impaired kidney function. Initial carboplatin dosages in these patients should be appropriately reduced (see DOSAGE AND ADMINISTRATION) and blood counts should be carefully monitored between courses. The use of carboplatin in combination with other bone marrow suppressing

therapies must be carefully managed with respect to dosage and timing in order to minimize additive effects.

Carboplatin has limited nephrotoxic potential, but concomitant treatment with aminoglycosides has resulted in increased renal and/or audiologic toxicity, and caution must be exercised when a patient receives both drugs. Clinically significant hearing loss has been reported to occur in pediatric patients when carboplatin was administered at higher than recommended doses in combination with other ototoxic agents.

Carboplatin can induce emesis, which can be more severe in patients previously receiving emetogenic therapy. The incidence and intensity of emesis have been reduced by using premedication with antiemetics. Although no conclusive efficacy data exist with the following chedules of carboplatin, lengthening the duration of single intravenous administration to 24 hours or dividing the total dose over five consecutive daily pulse doses has resulted in

Although peripheral neurotoxicity is infrequent, its incidence is increased in patients older than 65 years and in patients previously treated with cisplatin. Pre-existing cisplatin induced neurotoxicity does not worsen in about 70% of the patients receiving carboplatin as secondary treatment.

Loss of vision, which can be complete for light and colors, has been reported after the use of carboplatin with doses higher than those recommended in the package insert. Vision appears to recover totally or to a significant extent within weeks of stopping these high doses.

As in the case of other platinum-coordination compounds, allergic reactions to carboplatin have been reported. These may occur within minutes of administration and should be managed with appropriate supportive therapy. There is increased risk of allergic CONTRAINDICATIONS and ADVERSE REACTIONS, Allergic Reactions).

High dosages of carboplatin (more than 4 times the recommended dose) have resulted in severe abnormalities of liver function tests.

Carboplatin injection may cause fetal harm when administered to a pregnant woman Carboplatin has been shown to be embryotoxic and teratogenic in rats. There are no adequate and well-controlled studies in pregnant women. If this drug is used during pregnancy, or if the patient becomes pregnant while receiving this drug, the patient should be apprised of the potential hazard to the fetus. Women of childbearing potential should be advised to avoid becoming pregnant.

PRECAUTIONS

Needles or intravenous administration sets containing aluminum parts that may come in contact with carboplatin injection should not be used for the preparation or administration of the drug. Aluminum can react with carboplatin causing precipitate nation and loss of potency.

The renal effects of nephrotoxic compounds may be potentiated by carboplatin.

Carcinogenesis, Mutagenesis, Impairment of Fertility

The carcinogenic potential of carboplatin has not been studied, but compounds with similar mechanisms of action and mutagenicity profiles have been reported to be carcinogenic Carboplatin has been shown to be mutagenic both in vitro and in vivo. It has also been shown to be embryotoxic and teratogenic in rats receiving the drug during organogenesis.

Pregnancy Category D

It is not known whether carboplatin is excreted in human milk. Because there is a possibility of xicity in nursing infants secondary to carboplatin treatment of the mother, it is recommended that breast feeding be discontinued if the mother is treated with carboplatin injection.

Pediatric Use

Safety and effectiveness in pediatric patients have not been established (see WARNINGS,"audiologic toxicity").

Geriatric Use

Of the 789 patients in initial treatment combination therapy studies (NCIC and SWOG), 395

量は二

or other

Arial - 7 pt Patient Information: Arial - 10 pt

Last modified: 18 June 2015 11:55 AM

patients were treated with carboplatin in combination with cyclophosphamide. Of these, 141 were over 65 years of age and 22 were 75 years or older. In these trials, age was not a prognostic factor for survival. In terms of safety, elderly patients treated with carboplatin were more likely to develop severe thrombocytopenia than younger patients. In a combined database of 1,942 patients (414 were ≥ 65 years of age) that received single agent carboplatin for different tumor types, a similar incidence of adverse events was seen in patients 65 years and older and in patients less than 65. Other reported clinical experience has not identified differences in responses between elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out. Because renal function is often decreased in the elderly, renal function should be considered in the selection of carboplatin dosage (see DOSAGE AND ADMINISTRATION).

For a comparison of toxicities when carboplatin or cisplatin was given in combination with cyclophosphamide, see CLINICAL STUDIES, Use with Cyclophosphamide for Initial Treatment of Ovarian Cancer, Comparative Toxicity.

AD	VERSE EXPERIENCES IN	N PA	TIENTS WITH O	VARIAN CANCER
			First Line	Second Line
			Combination Therapy*	Single Agent Thera Percent
			Percent	
Bone Marrow				
Thrombocytope	nia < 100,000/mm³		66	62
	< 50,000/mm ³		33	35
Neutropenia	< 2,000 cells/mm ³		96	67
	< 1,000 cells/mm ³		82	21
Leukopenia	< 4,000 cells/mm ³		97	85
	< 2,000 cells/mm ³		71	26
Anemia	< 11 g/dL		90	90
	< 8 g/dL		14	21
Infections			16	5
Bleeding			8	5
Transfusions			35	44
Gastrointestinal				
Nausea and vor	miting		93	92
Vomiting			83	81
Other GI side ef	fects		46	21
Neurologic				
Peripheral neuro	opathies		15	6
Ototoxicity			12	1
Other sensory s	ide effects		5	1
Central neuroto	xicity		26	5
Renal				
Serum creatinin	e elevations		6	10
Blood urea elev	ations		17	22
Hepatic				
Bilirubin elevation	ons		5	5
SGOT elevation	s		20	19
Alkaline phosph	atase elevations		29	37
Electrolytes loss	S			
Sodium			10	47
Potassium			16	28
Calcium			16	31
Magnesium			61	43
Other side effec	ts			
Pain			44	23
Asthenia			41	11
Cardiovascular			19	6
Respiratory			10	6
Allergic			11	2
Genitourinary			10	2
Alopecia			49	2
Mucositis			8	1

'Use with Cyclophosphamide for Initial Treatment of Ovarian Cancer: Data are based on the experience

of 393 patients with ovarian cancer (regardless of baseline status) who received initial combination therapy with carboplatin and cyclophosphamide in two randomized controlled studies conducted by SWOG and NCIC (see **CLINICAL STUDIES**).

Combination with cyclophosphamide as well as duration of treatment may be responsible for the differences that can be noted in the adverse experience table.

†Single Agent Use for the Secondary Treatment of Ovarian Cancer: Data are based on the experience of 553 patients with previously treated ovarian carcinoma (regardless of baseline status) who received single

In the narrative section that follows, the incidences of adverse events are based on data from $1,\!893\,\text{patients with various types of tumors who received carboplatin as single agent the rapy.}$

Bone marrow suppression is the dose-limiting toxicity of carboplatin. Thrombocytopenia with et counts below 50,000/mm³ occurs in 25% of the patients (35% of pretreated ovariar cancer patients); neutropenia with granulocyte counts below 1,000/mm³ occurs in 16% of the patients (21% of pretreated ovarian cancer patients); leukopenia with WBC counts below 2.000/mm³ occurs in 15% of the patients (26% of pretreated ovarian cancer patients). The nadir usually occurs about day 21 in patients receiving single-agent therapy. By day 28, 90% of patients have platelet counts above 100,000/mm³; 74% have neutrophil counts above 2,000/mm³; 67% have leukocyte counts above 4,000/mm³

Marrow suppression is usually more severe in patients with impaired kidney function Patients with poor performance status have also experienced a higher incidence of severe leukopenia and thrombocytopenia.

The hematologic effects, although usually reversible, have resulted in infectious or hemorrhagic complications in 5% of the patients treated with carboplatin, with drug related death occurring in less than 1% of the patients. Fever has also been reported in patients

Anemia with hemoglobin less than 11 q/dL has been observed in 71% of the patients who started therapy with a baseline above that value. The incidence of anemia increases with increasing exposure to carboplatin. Transfusions have been administered to 26% of the patients treated with carboplatin (44% of previously treated ovarian cancer patients).

marrow suppressing drugs or with radiotherapy.

Vomiting occurs in 65% of the patients (81% of previously treated ovarian cancer patients) and in about one-third of these patients it is severe. Carboplatin, as a single agent or in mbination, is significantly less emetogenic than cisplatin; however, patients previously treated with emetogenic agents, especially cisplatin, appear to be more prone to vomiting. Nausea alone occurs in an additional 10% to 15% of patients. Both nausea and vomiting usually cease within 24 hours of treatment and are often responsive to antiemetic measures Although no conclusive efficacy data exist with the following schedules, prolonged administration of carboplatin, either by continuous 24-hour infusion or by daily pulse doses given for 5 consecutive days, was associated with less severe vomiting than the single-dose ntermittent schedule. Emesis was increased when carboplatin was used in combination with other emetogenic compounds. Other gastrointestinal effects observed frequently were pain, in 17% of the patients; diarrhea, in 6%; and constipation, also in 6%.

Neurologic Toxicity

Peripheral neuropathies have been observed in 4% of the patients receiving carboplatin (6% of pretreated ovarian cancer patients) with mild paresthesias occurring most frequently. Carboplatin therapy produces significantly fewer and less severe neurologic side effects than does therapy with cisplatin. However, patients older than 65 years and/or previously treated with cisplatin appear to have an increased risk (10%) for peripheral neuropathies In 70% of the patients with pre-existing cisplatin-induced peripheral neurotoxicity, there was no worsening of symptoms during therapy with carboplatin. Clinical ototoxicity and other ensory abnormalities such as visual disturbances and change in taste have been reported in only 1% of the patients. Central nervous system symptoms have been reported in 5% of the patients and appear to be most often related to the use of antiemetics.

Although the overall incidence of peripheral neurologic side effects induced by carboplatin is low, prolonged treatment, particularly in cisplatin pretreated patients, may result in cumulative

Nephrotoxicity

Development of abnormal renal function test results is uncommon, despite the fact that carboplatin, unlike cisplatin, has usually been administered without high-volume fluid hydration and/or forced diuresis. The incidences of abnormal renal function tests reported are 6% for serum creatinine and 14% for blood urea nitrogen (10% and 22%, respectively

in pretreated ovarian cancer patients). Most of these reported abnormalities have been mild and about one-half of them were reversible

Creatinine clearance has proven to be the most sensitive measure of kidney function in patients receiving carboplatin, and it appears to be the most useful test for correlating drug clearance and bone marrow suppression. Twenty-seven percent of the patients who had a baseline value of 60 mL/min or more demonstrated a reduction below this value during carboplatin therapy.

The incidences of abnormal liver function tests in patients with normal baseline values were reported as follows: total bilirubin, 5%; SGOT, 15%; and alkaline phosphatase, 24%; (5%, %, and 37%, respectively, in pretreated ovarian cancer patients). These abnormalities have generally been mild and reversible in about one-half of the cases, although the role of metastatic tumor in the liver may complicate the assessment in many patients. In a limited series of patients receiving very high dosages of carboplatin and autologous bone marrow transplantation, severe abnormalities of liver function tests were reported.

The incidences of abnormally decreased serum electrolyte values reported were as follows: sodium, 29%; potassium, 20%; calcium, 22%; and magnesium, 29%; (47%, 28%, 31%, and 43%, respectively, in pretreated ovarian cancer patients). Electrolyte supplementation was not routinely administered concomitantly with carboplatin, and these electrolyte abnormalities were rarely associated with symptoms.

Allergic Reactions

reactions have been similar in nature and severity to those reported with other platinumcontaining compounds, i.e., rash, urticaria, erythema, pruritus, and rarely bronchospasm and hypotension. Anaphylactic reactions have been reported as part of postmarketing surveillance (see WARNINGS). These reactions have been successfully managed with standard epinephrine, corticosteroid, and antihistamine therapy.

Injection Site Reactions

Injection site reactions, including redness, swelling, and pain, have been reported during postmarketing surveillance. Necrosis associated with extravasation has also been reported.

Other Events

their relationship to the tumor and to anemia was likely. Alopecia was reported (3%). Cardiovascular, respiratory, genitourinary, and mucosal side effects have occurred in 6% or less of the patients. Cardiovascular events (cardiac failure, embolism, cerebrovascular accidents) were fatal in less than 1% of the patients and did not appear to be related to chemotherapy. Cancer-associated hemolytic uremic syndrome has been reported rarely.

Malaise, anorexia, hypertension, dehydration, and stomatitis have been reported as part of postmarketing surveillance.

OVERDOSAGE

There is no known antidote for carboplatin injection overdosage. The anticipated complications of overdosage would be secondary to bone marrow suppression and/or hepatic toxicity.

DOSAGE AND ADMINISTRATION

NOTE: Aluminum reacts with carboplatin causing precipitate formation and loss of potency, therefore, needles or intravenous sets containing aluminum parts that may come in contact with the drug must not be used for the preparation or administration

Single Agent Therapy

Carboplatin injection, as a single agent, has been shown to be effective in patients with recurrent ovarian carcinoma at a dosage of 360 mg/m² IV on day 1 every 4 weeks (alternatively see Formula Dosing). In general, however, single intermittent courses of carboplatin should not be repeated until the neutrophil count is at least 2,000 and the platelet count is at least 100.000.

Combination Therapy with Cyclophosphamide

In the chemotherapy of advanced ovarian cancer, an effective combination for previously untreated patients consists of: Carboplatin - 300 mg/m² IV on day 1 every 4 weeks for 6 cycles (alternatively see Formula

Cyclophosphamide - 600 mg/m² IV on day 1 every 4 weeks for 6 cycles. For directions ding the use and administration of cyclophosphamide, please refer to its package insert (see CLINICAL STUDIES).

Intermittent courses of carboplatin in combination with cyclophosphamide should not be repeated until the neutrophil count is at least 2,000 and the platelet count is at least 100,000.

Dose Adjustment Recommendations

Pretreatment platelet count and performance status are important prognostic factors for severity of myelosuppression in previously treated patients.

The suggested dose adjustments for single agent or combination therapy shown in the table below are modified from controlled trials in previously treated and untreated patients with ovarian carcinoma. Blood counts were done weekly, and the recommendations are based on

Platelets	Neutrophils	Adjusted Dose * (From Prior Course)
> 100,000	> 2,000	125%
50 to 100,000	500 to 2,000	No Adjustment
< 50,000	< 500	75%

Percentages apply to carboplatin injection as a single agent or to both carboplatin and cyclophosphamide in combination. In the controlled studies, dosages were also adjusted at a lower level (50% to 60%) for severe myelosuppression. Escalations above 125% were not recommended for these studies.

Carboplatin injection is usually administered by an infusion lasting 15 minutes or longer. No pre- or post-treatment hydration or forced diuresis is required.

Patients with Impaired Kidney Function

Patients with creatinine clearance values below 60 mL/min are at increased risk of severe bone marrow suppression. In renally-impaired patients who received single agent carboplatin therapy, the incidence of severe leukopenia, neutropenia, or thrombocytopenia has been about 25% when the dosage modifications in the table below have been used.

25% When the decage meanications in the table below have been deca.		
Baseline Creatinine	Recommended	
Clearance	Dose on Day 1	
41 to 59 mL/min	250 mg/m ²	
16 to 40 mL/min	200 mg/m ²	

The data available for patients with severely impaired kidney function (creatinine clearance below 15 mL/min) are too limited to permit a recommendation for treatment.

These dosing recommendations apply to the initial course of treatment. Subsequent dosages should be adjusted according to the patient's tolerance based on the degree of bone marrow suppression.

Formula Dosing

Another approach for determining the initial dose of carboplatin injection is the use of mathematical formulae, which are based on a patient's pre-existing renal function or renal function and desired platelet nadir. Renal excretion is the major route of elimination for carboplatin (see CLINICAL PHARMACOLOGY). The use of dosing formulae, as compared to empirical dose calculation based on body surface area, allows compensation for patient variations in pretreatment renal function that might otherwise result in either underdosing (in patients with above average renal function) or overdosing (in patients with impaired renal

A simple formula for calculating dosage, based upon a patient's glomerular filtration rate (GFR in mL/min) and carboplatin injection target area under the concentration versus time curve (AUC in mg/mL•min), has been proposed by Calvert. In these studies, GFR was measured by ⁵¹Cr-EDTA clearance.

CALVERT FORMULA FOR CARBOPLATIN DOSING Total Dose (mg)=(target AUC) x (GFR + 25)

Note: With the Calvert formula, the total dose of carboplatin is calculated in mg, <u>not</u>

The target AUC of 4 mg/mL•min to 6 mg/mL•min using single agent carboplatin appears to provide the most appropriate dose range in previously treated patients. This study also showed a trend between the AUC of single agent carboplatin administered to previously treated patients and the likelihood of developing toxicity.

	% Actual Toxicity in Previously Treated Patients	
AUC (mg/mL•min)	Gr 3 or Gr 4	Gr 3 or Gr 4
	Thrombocytopenia	Leukopenia
4 to 5	16%	13%
6 to 7	33%	34%

Geriatric Dosing

Because renal function is often decreased in elderly patients, formula dosing of carboplating injection based on estimates of GFR should be used in elderly patients to provide predictable plasma carboplatin AUCs and thereby minimize the risk of toxicity

PREPARATION OF INTRAVENOUS SOLUTIONS

Carboplatin injection is a premixed aqueous solution of 10 mg/mL carboplatin

Carboplatin aqueous solution can be further diluted to concentrations as low as 0.5 mg/mL with 5% Dextrose in Water (D₅W) or 0.9% Sodium Chloride Injection, USP.

When prepared as directed, carboplatin aqueous solutions are stable for 8 hours at room temperature (25°C). Since no antibacterial preservative is contained in the formulation, it is recommended that carboplatin aqueous solutions be discarded 8 hours after dilution.

Product No.	NDC No.	
107205	63323-172-05	CARBOplatin Injection, 50 mg per 5 mL (10 mg per mL), in a 6 mL multiple-dose vial packaged individually.
107215	63323-172-15	CARBOplatin Injection, 150 mg per 15 mL (10 mg per mL), in a 20 mL multiple-dose vial packaged individually.
107245	63323-172-45	CARBOplatin Injection, 450 mg per 45 mL (10 mg per mL), in a 50 mL multiple-dose vial packaged individually.

The container closure is not made with natural rubber latex

STORAGE

Unopened vials of carboplatin injection are stable to the date indicated on the package when stored at 25°C (77°F); excursions permitted from 15° to 30°C (59° to 86°F) [see USP

PROTECT FROM LIGHT.

Carboplatin injection multidose vials maintain microbial, chemical, and physical stability for up to 14 days at 25°C following multiple needle entries.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration. Solutions for infusion should be discarded 8 hours after preparation.

HANDLING AND DISPOSAL

Caution should be exercised in handling and preparing carboplatin injection. Several guidelines on this subject have been published1-4

To minimize the risk of dermal exposure, always wear impervious gloves when handling vials containing carboplatin injection. If carboplatin injection contacts the skin, immediately wash the skin thoroughly with soap and water. If carboplatin injection contacts mucous membranes, the membranes should be flushed immediately and thoroughly with water. More information is available in the references listed below.

- 1. NIOSH Alert: Preventing occupational exposures to antineoplastic and other hazardous drugs in healthcare settings. 2004. U.S. Department of Health and Human Services. Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 2004-165.
- 2. OSHA Technical Manual, TED 1-0.15A, Section VI: Chapter 2. Controlling occupational exposure to hazardous drugs. OSHA, 1999. http://www.osha.gov/dts/osta/otm/otm_vi/ otm vi 2.html.
- 3. American Society of Health-System Pharmacists. ASHP guidelines on handling hazardous drugs. Am J Health-Syst Pharm. 2006: 63:1172-1193.
- 4. Polovich M, White JM, Kelleher LO, eds. 2005. Chemotherapy and biotherapy guidelines and recommendations for practice. 2nd ed. Pittsburgh, PA: Oncology Nursing Society.

Manufactured for FRESENIUS KABI

Fresenius Kabi USA, LLC

Made in India

For Product Inquiry: 1-800-551-7176 or www.fresenius-kabi.us

451167B/Revised: June 2015

7220132221



羉 A20132221 USA САВОРСАТІИ ІИЈЕСТІОИ

> Manufactured for: FRESENIUS KABI Fresenius Kabi USA, LLC

Lake Zurich, IL 60047 451167B/Revised: June 2015

y and liver problem:
of hearing or ringin
ears;

is is

Arial - 7 pt Patient Information: Arial - 10 pt Ver. 11

Last modified: 18 June 2015 11:55 AM